

Forest structure comparison of Hutcheson Memorial Forest and eight old woods on the New Jersey Piedmont

Richard T.T. Forman and Bruce A. Elstrom, Department of Botany, Rutgers University

Forman, Richard T. T., and Bruce A. Elstrom. (Dep. Botany, Rutgers Univ., New Brunswick, N.J. 08903). Forest structure comparison of Hutcheson Memorial Forest and eight old woods on the New Jersey Piedmont. Hutcheson Mem. Forest Bull. 3:44-51, 1975.—Compared with 8 nearby oak woods, the William L. Hutcheson Memorial Forest is higher in number of tree species, percent exotic species, percent understory, subcanopy and large shrub layer species, seedlings of *Fraxinus americana*, *Acer platanoides* and the red oak group, saplings of *Fraxinus americana*, trees of *Cornus florida* and *Quercus alba*, and range of soil moisture conditions. HMF is typical in the number of soil types present and the percents of canopy, edge, central closed canopy, and lowland species. It is lower than usual in seedlings, saplings and trees of *Acer rubrum*, *A. saccharum* and *Fagus grandifolia*, seedlings of *Prunus serotina*, and trees of *Quercus velutina*. Relative to the other woods, at HMF *Fraxinus americana* and *Acer rubrum* are likely to continue their unusual importances, *Cornus florida* is likely to become more typical, and *Acer platanoides* and *Prunus serotina* more atypical. Relative to the present HMF structure *Acer saccharum* is likely to increase in importance, and *Quercus alba*, *Carya* spp. and *Prunus avium* decrease. The higher tree species diversity at HMF, characterized by species of lower forest strata and many exotics, may be related to higher people use. Community coefficient values and composition of dominant species indicate Hutcheson Memorial Forest is broadly representative of the forests on the upland red shale New Jersey Piedmont.

An exceptional amount of research has been done at the William L. Hutcheson Memorial Forest in the last 25 years, with over 80 published papers and reports based on work in the woods and adjacent fields (Small 1973). The forest structure has been described as a mixed *Quercus* canopy, *Cornus florida* understory, and *Viburnum* shrub layer (Bard 1952; Buell 1957; Monk 1957, 1961a; Sulser 1971). Vascular plant species diversity includes 40 trees, 39 shrubs and 232 herbs (Frei and Fairbrothers 1963). Fifteen percent of the tree species and 22% of the total species are exotic (alien) species, some of which are increasing in dominance (Frei and Fairbrothers 1963; Ambler 1965; Sulser 1971). The species abundances of the forest edge are relatively distinct from the closed canopy, which, however, contains numerous gaps (Wales 1972). Though this old forest has apparently been unburned for 3 centuries and uncut, major windstorms and perhaps pest outbreaks have caused pulses of change in the woods, and it is suggested that maples, beech and ash are slowly increasing at the expense of oaks and hickories (Buell 1957; Monk 1961b; Reiners and Reiners 1965; Sulser 1971). Are these forest characteristics and changes representative of the woods on the red shale Piedmont of New Jersey, or is this an atypical woods for the region?

The region is broadly classified as oak-chestnut forest in a study of the eastern United States (Braun 1950) or

mixed oak forest in a study of New Jersey (Robichaux and Buell 1973). Detailed studies of forest structure: Cushtunk Mountain (Cantlon 1953), Voorhees State Park (McDonough and Buell 1956), Watchung Reservation in Union Co. (Baird 1956), Herrontown Woods Princeton (Kramer 1971), and the Institute Woods Princeton (Horn 1971, 1975) are all on distinct geologic substrata and soils adjacent to the red shale Piedmont. Lowland woodlands in the Piedmont are described for the Raritan River (Buell and Wistendahl 1955; Wistendahl 1958), Millstone River (VanVechten and Buell 1959) and Stony Brook (Horn 1971, 1975). In the apparent absence of detailed studies in other woods of the upland, red shale Piedmont, Hutcheson Memorial Forest (HMF) has been assumed to be representative of this region.

The objectives of this study are to (a) compare the Hutcheson Memorial Forest with other woods for numbers, types, composition and importances of tree species, and (b) look for evidence of relative future changes between HMF and the other woods, as well as within Hutcheson Memorial Forest itself.

Methods

Eight woods surrounded by fields, in addition to Hutcheson Memorial Forest, were selected in a rural New Jersey area approximately 32 km in diameter and bounded by New Brunswick, Oldwick, Flemington and Princeton. The woods are 20-100 m elevation in a Triassic red shale region of the Piedmont (Lewis and Kummel 1915; Widmer 1964). Soils are predominantly silt and shaly loams of the Penn (including Norton) series of the Brunswick formation (Tedrow 1963; Ugolini 1964; U.S. Soil Conservation Service 1958-1967). The climate is mild, with approximately 100 cm precipitation relatively evenly distributed through the year, average annual temperature 11.7°C, and an average monthly temperature range of 0.0-24.0°C (Biel 1958; U.S. Weather Bureau 1959).

Criteria for woods selection were the presence of mature trees (60-100 cm dbh, diameter measured 1.5 m above ground level) over the entire woods, all forest structural layers (canopy, subcanopy, understory, shrub and herb), discreteness from the surrounding fields, approximately isodiametric shape, mature edges, large size, slope not exceeding 10°, well drained soil, lack of disturbance (recent fire, cutting, etc.), absence of streams, and distance from population centers and pollution sources. Three size classes were used, 7.5 ha, 10 ha and 24 ha, and one woods of each size was selected in the

northern portion of the study area, one of each size in the central portion, and one of each size in the southern portion. Hutcheson Memorial Forest is the 24 ha woods in the central portion. In the following descriptions of the woods, location and soils are given, plus any exceptions to the above listed criteria for woods selection.

- N7.5.* (7.5 ha woods in the northern portion), Maryknoll Farm West. Three km north of Whitehouse Station, approximately 0.6 km west of route 523. Soils: 60% Norton loam eroded, 40% Norton loam. Gaps common in portion due to recent cutting, edge indistinct in places.
- C7.5.* Wohendorf-Perry Woods. Three km south of Whitehouse Station, at end of Edgewood Rd., 1 km east of route 523. Soils: 45% Penn shaly silt loam, 45% Penn shaly silt loam eroded, 10% Reaville silt loam wet. Rectangular shape, temporary stream.
- S7.5.* Water Tower Woods. Northeast of Blawenburg, approximately 3 km north of route 518, 2 km west of route 206, north side of Sunset Road. Soils: 70% Royce silt loam, 5% Royce silt loam eroded, 10% Lansdowne silt loam, 15% Birdsboro gravelly loam. Understory with non-uniform coverage, some wood cut for firewood, large water tower.
- N10.* Maryknoll Farm East. Three km north of Whitehouse Station, approximately 0.3 km west of route 523. Soils: 80% Norton loam eroded, 20% Norton loam. Gaps common in portion due to recent cutting, edge indistinct in places.
- C10.* Reno Farm. South of Whitehouse Station, approximately 2 km north of Pleasant Run, by high tension wires on east side of Cole Road. Soils: 40% Penn shaly silt loam eroded, 40% Reaville silt loam, 20% Penn-Bucks complex. Rectangular in shape, permanent stream, understory with non-uniform coverage, 1972 ground fire in one corner.
- S10.* Ringoes Corso Woods. Three km west of Ringoes, approximately 0.3 km south of Sergeantsville Rd. Soils: 90% Penn shaly silt loam eroded, 10% Rough broken land, shale (bedrock at surface with patches of thin soil). Permanent stream, edge indistinct in places, house near edge.
- N24.* Treetops. Two km south of Lamington, south of interstate highway 78, west of Lamington River, and immediately west of Fiddlers Elbow golf course. Soils: 40% Athol gravelly loam, 20% Lansdowne silt loam, 20% Annandale and Edneyville gravelly loam, 10% Athol gravelly loam eroded, 5% Reaville silt loam wet, 5% Norton loam eroded. Standing water, gaps common in portion due to recent cutting, houses near edge, soils mainly developed from glacial till and calcareous conglomerate.
- C24.* William L. Hutcheson Memorial Forest. One km east of East Millstone, on south side of route 514. Soils: 50% Penn shaly silt loam, 20% Royce silt loam, 15% Croton silt loam, 15% Lansdowne silt loam. Rectangular shape, permanent stream,

standing water, gaps common due to blowdowns, western portion beyond the narrows excluded.

- S24.* Western Electric Woods. One km south of Mt. Rose and 1 km west of route 569. Soils: 70% Reaville silt loam, 8% Penn shaly silt loam, 5% Bowmansville silt loam, 5% Klinesville shaly loam, 5% Doylestown silt loam, 3% Reaville silt loam wet, 3% Reaville silt loam eroded. Permanent stream, standing water, understory non-uniform coverage, and grazed, burned or cleared recently near some edges.

Sampling for tree species was done between August 1972 and September 1973, and 3 methods were utilized: quadrats, transects and reconnaissance. A series of parallel lines, marked with colorful plastic flagging, divided each woods into 30 m wide sections, running perpendicular to the straightest long edge of the woods. Quadrats were located in a stratified random manner throughout the woods (including the edge portions) using the marked sections. Quadrats were 40 x 10 m for trees (≥ 10 cm dbh), 10 x 2.5 m for saplings (< 10 cm and ≥ 2.5 cm dbh), and 2.5 x 0.5 m for seedlings (< 2.5 cm dbh). The sapling and seedling quadrats were nested in a constant corner of the tree quadrats. All individuals exhibiting a single-stemmed growth pattern, excluding vines, and a dbh ≥ 2.5 cm were included as trees or saplings. All seedlings of those species which at least occasionally attain tree or sapling status in the study area were included. Hutcheson Memorial Forest was sampled with 116 quadrats. *N10* woods with 55, *S7.5* woods with 45, and based on the point where species-area curves for these 3 woods begin to level off, 10 quadrats were done in the remaining woods. Presence and abundance of species were recorded.

Transects were 10 x 90 m running perpendicular to the edge of the woods and from the outer canopy level limbs toward the center of the woods. Transects were located randomly along the straightest east or west edge, but no closer than 30 m from a north or south edge. Presence, abundance and dbh of trees and saplings were recorded. Fourteen transects were sampled at the *N10* woods, 8 at *S7.5*, 4 at *C24* (HMF), and 3 at the remaining woods. In the absence of good east and west edges with well drained woods at Hutcheson Memorial Forest, 2 transects were from the south edge between the trail and the "dog leg" and 2 from the north edge opposite these (see map in Monk 1957).

Reconnaissance was done by walking at approximately 10 m intervals in a zig-zag fashion between the marked lines 30 m apart. Species absent from the quadrats and transects were recorded. Distance traversed and time in reconnaissance sampling were approximately proportional to the area of the woods.

Tree species were identified with Fernald (1950), Gleason (1962), Harlow (1959) and the Chrysler Herbarium at Rutgers University. We divided the species into 4 growth form categories based on the normal maximum level attained in the forests: canopy, subcanopy, understory and large shrub. Species were classified as edge species, closed canopy species, and lowland species

according to Wales (1969), Agriculture Handbook No. 271 (1965), Harlow (1959), and observations of distributions in this study. Species were classified as exotic if not native east of the Appalachian Mountains. More detailed descriptions of study areas, tree and soil sampling are given in Galli (1973) and Elfstrom (M.S. thesis in preparation).

Soil types were determined from detailed maps of the soils in each county (U.S. Soil Conservation Service 1958-1967). All samples for soil moisture were collected on the same day in late May 1973, weighed, dried at 105°C for 48 hrs, and reweighed to determine percent weight loss. Ten samples were taken in each woods. If there were no obviously wet or dry areas, samples were taken in a stratified random manner using the marked lines 30 m apart. When distinctly wetter or drier areas were evident in a woods, 5 samples were taken in the wetter areas and 5 samples in the drier areas, in order to determine the range of soil moisture conditions present.

Results

We encountered a total of 41 species with a dbh ≥ 2.5 cm in Hutcheson Memorial Forest based on quadrat sampling and reconnaissance (table 1). The average total number of species in the other 8 woods was 32.4 (standard deviation = 3.0), or if only the two 24 ha woods equal in size to HMF are considered, 34.0 (sd = 5.7). Therefore species diversity, measured as number of species, appears higher at Hutcheson Memorial Forest than at the other woods in the region. When the quadrat data alone are compared (table 1), based on 10 quadrats selected in a stratified random fashion in all woods, we find: HMF 16 species, 8-woods average 16.9 (sd = 1.3), and 2-woods average 16.5 (sd = 0.7). Thus the greater species diversity at Hutcheson Memorial Forest is due to rare species not encountered in 10 stratified random quadrats.

In examining this difference, first we will consider the types of species and later the particular species present. A notable 14.6% of the total species encountered at HMF are exotic species (table 1). This compares with the 8-woods average of 6.1% (sd = 2.1%). Of the 6 exotic species at HMF, 2 may be considered common, while the average 2.0 (sd = 0.8) exotic species in the other woods are virtually always rare. Thus exotic species are more prevalent at HMF than usual in the region.

Comparing vertical strata in the woods we found 12 canopy species at Hutcheson Memorial Forest (table 1) vs the 8-woods average of 13.5 (sd = 1.2) or the 2-woods average of 14.5 (sd = 0.7), indicating little difference, or perhaps a slightly lower canopy diversity at HMF. There were 11 subcanopy species at HMF vs 7.6 (sd = 2.1) average in the 8 woods or 8.5 (sd = 3.5) in the 2 woods, suggesting a significant difference. Eight understory species at HMF compared with the 8-woods average 5.6 (sd = 0.5) or the 2-woods average 5.0 (sd = 0.0), a notable difference. In the large shrub category HMF had 8 spe-

cies while the 8-woods average was 5.8 (sd = 0.9) or the 2-woods average 6.0 (sd = 1.4), again a notable difference. Therefore the greater species diversity at HMF is due to the greater number of species in the lower strata of the forest. It is notable also that the diversity of HMF canopy is not greater than the other woods.

Some species have distinctly higher relative abundances near the edge of a forest, some in the center, some in the closed canopy portion, and some are evenly distributed from the edge to the center. At HMF we found 32% of the total species were edge species while the 8-woods average was 31.5% (sd = 4.3); the pattern was similar in the quadrats only. Closed canopy species at HMF were 24.4% of the total vs 29.5% (sd = 3.9) in the other woods, with little difference between the two quadrats. These results indicate that the percents of edge and closed canopy species are typical at Hutcheson Memorial Forest.

Lowland species, primarily in poorly drained spots along infrequent streams, comprise 24.4% of the total species at HMF and an average 21.2% (sd = 3.0) in the other woods. The quadrat samples gave 16.0% in HMF and 17.6% (sd = 7.1) respectively. Thus lowland species at HMF are of about equal prevalence as in the other woods.

To compare the relative similarity of the total species present at HMF with the other woods we calculate community coefficients (CC) for pairs of woods (Braun and Curtis 1957):

$$CC = \frac{2 S_c}{S_1 + S_2}$$

S_1 is the number of species at HMF, S_2 the number of species in a second woods, and S_c the number of species common to both woods. The CC's comparing HMF with each of the 8 woods were 0.73 (for the N7.5 woods), 0.73 (C7.5), 0.65 (S7.5), 0.76 (N10), 0.72 (C10), 0.82 (S10), 0.76 (N24) and 0.79 (S24), with an average CC of 0.73 (sd = 0.05). These are high values, considering replicate samples within such woods commonly have a CC of about 0.80-0.85, and indicates that the HMF tree community is quite similar overall to the other woods. The rather low variability indicates further that it is similar to essentially all of the woods.

Four species were encountered only at HMF (table 1): *Paulownia tomentosa*, *Acer negundo*, *Cornus racemosa*, and *Pyrus malus*. One is an exotic, 2 are subcanopy species, 1 understory and 1 large shrub species. In the other woods had an average of 1.3 (range 0.3) unique species, those being *Ulmus rubra*, *Tsuga canadensis*, *Robinia pseudo-acacia*, *Pinus virginiana*, *Platanus occidentalis*, *Populus tremuloides*, *Maclura pomifera*, *Prunus pennsylvanica*, *Rhus copallina* and *Populus grandidentata* were in two or more woods but not encountered at HMF. Of the 13 species only in the 8 woods, 1 is an exotic, 6 are canopy species, 3 subcanopy, 1 understory, and 3 large shrub species. Therefore, based on rare species either unique to HMF or unique to the other woods, there is a slightly higher number of such species

Species	N	N	N	N	N	N	N	N	N	N	N
<i>Penus pennsylvanicus</i> L. LS, ED	0	0	0	0	0	0	0	0	0	0	0
<i>Liriodendron tulipifera</i> L. LS, ED, EX	0	0	0	0	0	0	0	0	0	0	0.2
<i>Carya cordifolia</i> (Wang) K. Koch CA, CL, LO	R	0	0	0	0	R	0	0	0	0	0
<i>Carya alaba</i> (Mill.) Sweet CA, CL	R	R	0	R	R	R	R	R	R	R	R
<i>Carya ovata</i> (Wang) Surg. CA, CL	R	R	R	R	R	R	R	R	R	R	R
<i>Carya ovata</i> (Mill.) K. Koch CA, CL	R	R	R	R	R	R	R	R	R	R	R
<i>Carya tomentosa</i> Nutt. CA	0	0	R	R	R	R	0	0	0	0	0
<i>Panicum compositum</i> (Thunb.) Spreng. S, ED, EX	0	0	0	0	0	0	0	0	0	0	0
<i>Pinus virginiana</i> Mill. S, ED	0	0	0	0	0	R	0	0	0	0	0
<i>Platanus occidentalis</i> L. CA, LO	0	0	0	0	0	R	0	0	0	0	0
<i>Tsuga canadensis</i> (L.) Carv. CA, CL	0	0	0	R	0	0	0	0	0	0	0
<i>Rhus americana</i> L. CA, LO	0	R	0	R	R	R	R	R	R	R	R
<i>Vincetoxicum</i> Mill. CA, LO	0	0	0	0	0	0	0	R	0	0	0
<i>Berula populifolia</i> Marsh. S, ED	R	0	0	R	R	0	R	R	0	0	0
<i>Diospyros virginiana</i> L. S, ED, LO	0	0	0	0	0	0	R	0	0	0	R
<i>Populus grandidentata</i> Michx. S, ED	0	0	0	R	R	0	0	0	0	0	0
<i>Populus tremuloides</i> Michx. S, ED	0	0	0	0	0	0	R	0	0	0	0
<i>Sambucus pseudo-racemosa</i> L. S, ED	0	0	0	0	0	R	0	0	0	0	0
<i>Tilia americana</i> L. S, CL	0	0	0	0	0	R	0	0	0	0	0
<i>Corylus americana</i> Walt. LS, ED	0	0	R	0	0	0	R	0	0	0	R
<i>Cornus racemosa</i> Lam. LS	0	0	0	0	0	0	0	0	0	0	0
<i>Maclura pomifera</i> (Raf.) Schneid. U, ED, EX	0	0	R	0	0	0	0	0	0	0	0
<i>Carya virginiana</i> (Mill.) K. Koch U, CL	0	R	0	0	R	R	0	0	0	0	R
<i>Rhus copallina</i> L. LS, ED	R	0	0	0	0	0	0	0	0	0	0
<i>Rhus typhina</i> L. LS, ED	R	0	R	R	0	R	0	0	0	0	R
<i>Spiraea tomentosa</i> L. LS, ED	0	0	R	0	0	0	0	0	0	0	0

Some differences are seen when we compare the relative basal areas of species at HMF and the average for the 8 woods (table 2). (a) *Cornus florida* is much higher in basal area than in the other woods, (b) *Quercus alba* is somewhat higher, (c) *Quercus velutina*, *Carya* spp., *Fagus grandifolia*, *Acer saccharum*, and *Liriodendron tulipifera* are somewhat lower, and (d) *Acer rubrum* is much lower in basal area than in the others. Therefore, based on relative abundance and basal area of trees at HMF, 2 species, *Cornus florida* and *Quercus alba*, are more important than in the other woods, and at least 4 species, *Acer rubrum*, *Quercus velutina*, *Fagus grandifolia*, and *Acer saccharum*, are less important than usual.

The differences in successional status of species may be suggested by the relative abundances of saplings and seedlings (table 1). Sapling data at HMF show (a) *Fraxinus americana* is much more abundant than the

8-woods average, (b) *Acer rubrum* and *A. saccharum* somewhat less abundant, and (c) *Fagus grandifolia* much less abundant than the other woods. In contrast, seedling data at HMF (table 1) show the red oak group seedlings much more abundant than the other woods, (b) *Fraxinus americana* and *Acer platanoides* somewhat more abundant, (c) *Acer saccharum*, *Prunus serotina*, *Fagus grandifolia* and *Hamamelis virginiana* somewhat less abundant, and (d) *Acer rubrum* much less abundant than the 8-woods average.

We now examine the changes at Hutcheson Memorial Forest relative to the other woods, suggested by the tree, sapling and seedling results. *Fraxinus americana* which is presently more important at HMF than the 8-woods average in all 3 categories, may continue to maintain or increase its higher relative importance. The striking difference in the number of large *Cornus florida* trees and to a lesser extent *Quercus alba* at HMF vs the other woods may disappear through time, since the sapling and seedling relative abundances are similar. *Acer rubrum*, *A. saccharum*, and *Fagus grandifolia*, presently less important in all 3 categories at HMF than elsewhere, may continue at a lower level than in the other woods. The relatively low importance of *Quercus velutina* at HMF compared with the other woods may disappear.

Table 2. Relative basal areas of species in transects. Based on trees ≥ 10 cm dbh. Transects 10 x 90 m from edge toward center of woods. Relative basal areas in %.

Size of Woods (ha)	7.5	7.5	7.5	10	10	10	24	24	Average of 8 woods
Portion of Study Area (North, Central, South)	N	C	S	N	C	S	N	S	
Number of Transects	3	3	8	16	3	1	1	3	
Number of Trees	82	96	127	267	87	20	22	74	
Total Basal Area (cm^2)	2016	2379	3561	8075	1858	1664	1891	2195	
<i>Cornus florida</i>	3.6	2.3	1.8	10.4	0	13.9	9.8	0	6.0
<i>Quercus alba</i>	4.6	22.6	23.7	13.5	25.0	7.5	12.2	14.5	16.0
<i>Quercus rubra</i>	6.5	23.2	1.1	22.4	32.9	10.4	0	23.2	17.4
<i>Quercus velutina</i>	22.9	11.1	0.8	16.6	28.7	19.2	11.7	0	19.2
<i>Fraxinus americana</i>	17.1	11.0	0.8	0.2	0	16.3	0	5.3	5.0
<i>Prunus avium</i>	0	2.9	1.4	1.0	0	3.4	0	0.6	1.5
<i>Carya</i> spp.	2.0	11.7	1.2	10.9	6.5	3.5	5.1	8.7	11.1
<i>Acer platanoides</i>	0	0	0	0	0	0	0	0	0
<i>Sassafras albidum</i>	1.1	4.1	0	0.6	0	2.9	0	0.5	1.3
<i>Ulmus</i> spp.	0	0	0	0.3	0	0	0	0	0.0
<i>Acer rubrum</i>	14.9	3.3	11.4	10.6	8.7	1.9	25.6	11.4	12.8
<i>Fagus grandifolia</i>	17.2	0	1.3	10.4	0	0	10.8	6.5	6.6
<i>Acer saccharum</i>	0	0.9	0	0.6	0	30.8	2.3	13.4	5.0
<i>Liriodendron tulipifera</i>	7.4	0	0	0.3	0	0	22.9	1.8	6.7
<i>Quercus palestris</i>	5.4	0	0.7	1.1	0	0	0	4.5	1.7
<i>Quercus prinus</i>	4.7	0	0	0	0	1.8	0	2.7	1.3
<i>Juglans nigra</i>	4.2	0	0	0	0	1.1	0	0	0.8
<i>Prunus americana</i>	0.5	0	0	0.2	0	0.8	0	0	0.2
<i>Berula populifolia</i>	0.6	0	0	0	0	0	0	0	0
<i>Myrica sylvatica</i>	1.5	0	0.3	1.3	0	0	0	1.8	1.0
<i>Amelanchier arborea</i>	0.5	4.1	0	0	1.5	0	0	0	0.8
<i>Castanea dentata</i>	0	0	0	0.1	0	0	0	0	0.0
<i>Allanhus altissim</i>	0	0	0	0	0	0	0	0	0.0

Table 3. Soil type and soil moisture conditions in the woods. Soil types determined from county soil maps. Soil moisture in percent by weight based on 10 samples per woods with the wettest and driest areas in a woods sampled when evident. All samples taken in one day.

Size of Woods (ha)	1.5	7.5	7.5	10	10	10	25	25	Average of 9 woods (HMF)	24
Location of Study Area (North, Central, South)	A	C	S	N	C	S	N	S		C
Total Number of Species	31	29	30	32	31	35	38	10	32.5	41
Number of Soil Types	2	3	1	2	1	2	4	7	3.5	4
Maximum Soil Moisture %	32.3	32.2	37.0	31.3	30.0	34.0	33.3	32.6	30.3	33.9
Minimum Soil Moisture %	21.0	20.0	28.0	25.5	25.5	29.7	25.3	35.3	26.1	25.7
Range of Soil Moisture %	9.3	22.2	9.0	6.8	15.5	14.3	7.8	19.1	14.3	11.3

appear in time, if many of the red oak group seedlings are this species, but this remains uncertain. The seedling values for the red oak group and *Acer platanoides* at HMF are higher than in the other woods, such that these species may become relatively more important. Similarly *Prunus serotina* may become less important at HMF compared to elsewhere. Note particularly that these trends for HMF are relative to the other woods, i.e. in comparison with the trends in the 8-woods averages.

The suggested trends within HMF itself, based on comparing the tree data of tables 1 and 2 with the seedling-sapling data (table 1), of course are different. Here *Cornus florida*, *Fraxinus americana*, the red oak group (primarily *Quercus rubra* and *Q. velutina*), and the less common species may change little in relative importance. However, *Acer saccharum* may increase in relative importance, while *Quercus alba*, *Carya* spp. and *Prunus avium* may decrease within Hutcheson Memorial Forest.

Two environmental factors, soil type and soil moisture, were studied to see if they correlated with the higher species diversity at HMF than the other woods. The 4 soil types encountered at HMF (table 3) are essentially the same number as the average 3.5 (sd = 1.9) for the other 8 woods, and slightly less than the 2-woods average of 6.5 (sd = 0.7). If the number of soil types and the number of species per woods are compared for all 9 woods, there is a slight positive correlation, but with high variability. However, if the number of soil types and species per woods are compared within the 3 (woods) size groups separately (7.5 ha, 10 ha and 24 ha), in each case there is a slight negative correlation (indicative of a size-soil type interaction). We conclude therefore that the number of soil types at HMF and the other woods is unimportant in determining the difference in species diversity.

The range of soil moisture conditions (table 3) within HMF, 31.3%, exceeds the 8-woods average range of 14.3% (sd = 5.4) and the 2-woods range of 18.5% (sd = 0.9). Caution should be exercised in interpreting these results since soil moisture is measured as percent by weight, without measurements of soil texture. A plot of soil moisture range vs number of species for the 9 woods would show a positive correlation, but with an exceed-

ingly large variation. If these are plotted separately for the 3 (woods) size groups however, two of the correlations are negative and one positive, again with considerable variation. These results indicate that, though the range of soil moisture conditions appears greater at HMF than in the other woods, this is likely no more than a minor contributing factor to the higher species diversity at Hutcheson Memorial Forest.

Discussion

The results of tree, sapling, seedling and soil sampling indicate a higher species diversity, especially of species in lower forest strata (of which many are exotics), at Hutcheson Memorial Forest than the other woods. However, with the exception of a minor effect of soil moisture, the data give few clues on the reason for the higher diversity. Two possibilities require evaluation: sampling methods and unsampled factors.

Since the number of species in 10 quadrats at HMF (16 species) and the other woods (16.9 species, sd = 1.3) is the same, the diversity difference comes in the subsequent sampling. The number of species added after the 10 quadrats is: 25 at HMF based on 106 more quadrats plus reconnaissance, 15 at S7.5 based on 35 more quadrats plus reconnaissance, 17 at N10 based on 45 more quadrats plus reconnaissance, and an average 15.3 (sd = 3.6) in the remaining 6 woods based on reconnaissance alone. Thus in the other 8 woods the number of species added is about the same whether based on quadrats plus reconnaissance or reconnaissance alone, and is considerably less than at HMF. There is little difference in the number of added species in the 106 quadrats at HMF (9 species) and the 35 at S7.5 (8 species) or 45 at N10 (6 species), which combined with the previous point indicates that the large increase in diversity at HMF is not due to the intensive sampling of the 106 additional quadrats. We conclude that sampling differences between HMF and the other woods contribute little, if anything, to the higher diversity recorded at HMF.

The unsampled factors which seem most important to us are people use and proximity to population centers. People usage at HMF includes approximately 40-50 group tours annually which are limited to the 1.5 km path at the eastern end. A few dozen research projects, strictly limited usage by small advanced classes, and caretaking of the property bring people throughout the woods on a fairly regular basis. Though the numbers have varied, this usage has continued for approximately two decades. We know far less about people usage of the 8 woods, other than our own observations during year-round sampling in this and 6 associated studies. The major uses appear to be hunters during a few weeks in the fall, hikers infrequently, children infrequently, and at N24 horseback riders infrequently. People may carry seeds, presumably involuntarily, into the forest and the expected result would be in most cases, few individuals of tree species. Since people usage is believed greater in

the past 20 years than earlier, trees introduced by people would mostly be less than 20 years old. This hypothesis gibes with the observed results where most of the additional species at HMF are rare, composed of individuals <10 cm dbh and in the lower strata of the forest.

We note that limiting people usage is an important management priority at the William L. Hutcheson Memorial Forest.

The number of exotic species at HMF might be related to proximity to population centers where exotic species are of greatest importance. However, HMF is 11 km from New Brunswick and Somerville and the 8-woods average is 11.0 km (sd = 3.5) from the nearest major population center (New Brunswick, Somerville, Flemington or Princeton). Therefore distance from population centers does not correlate with the higher diversity at HMF. It might be noted that the large areal spread of housing developments around these population centers, where exotic plantings are abundant, has taken place in the last 2 decades.

The distance from small villages which originated as much as two centuries ago might also be examined, since HMF is only 1 km east of such a village, East Millstone. Distances from each of the 8 woods to its nearest village are 1, 2, 3, 3, 3, 3, 3, and 4 km. Therefore HMF is significantly closer to such a village than the 8-woods average, 2.8 km. This could be a contributing factor to the higher species diversity, but the importance of this is unknown.

The indicated changes for the future of the woods are based on the assumption that the relative proportions of species as seedlings and saplings will be important in determining relative proportions of future canopy species. Unfortunately we know too little about the effects of pulses of tree reproduction and growth, subtle climatic changes, air pollution, changes in herbivore populations, and fire history. Thus we must cautiously live with this assumption for predictions, and hope that studies of a few decades in the lifetime of trees are representative.

Though the actual species composition of the HMF canopy may change, the diversity may not, since the higher diversity at HMF is of species which only reach to lower strata. This assumes these species, including several exotics, do not prevent other species from reaching the canopy. Studies such as those of Horn (1971, 1975) showing differential reproduction under each tree species seem particularly valuable.

The disparate data for *Acer rubrum* at HMF in tables 1 and 2 are due to the relative absence of the species in the central portion of the forest where the transects were done. Table 1 is representative of *Acer rubrum* for the entire forest. We are uncertain of future trends in the red oak group in any woods because the seedlings were not separated by species. Also the red oak group is the only case where there may be a significant difference in the production of seedlings between 1972 and 1973 (table 1). Based on the relative abundance of trees, the seedlings are probably almost all *Quercus rubra* and *Q. velutina*. At HMF the two species are about equal in abundance, but in all of the other woods *Quercus velutina* is more abundant, and often many times more abundant, than

Q. rubra. Differential changes in the two species probable but unpredictable.

The future changes in species composition to be expected at HMF based on intensive sampling throughout the woods at one time, agree overall with Sulser's significant study (1971), based on sampling one portion comparing data with a similar sample 20 years earlier. Differences for some individual species are evident, expected since the present study includes a broader and >20% of the forest area.

The predicted increase in northern hardwoods species like *Acer saccharum* and *Fraxinus americana* at the expense of oak forest species like *Quercus alba* and *Q. spp.* in these 9 woods on the upland red shale Piedmont agrees also with results on other geologic substrata in Piedmont and to the North in New Jersey (Buell 1966). Such a change in forest structure over a broad area would have major ecological implications.

We appreciate the field assistance of A.E. Galli, M. Engelmann, E.W. Murray and V.A. Rudis, the cooperation of the land owners, and the financial assistance of the U.S. Forest Service Pinchot Institute Consortium for Environmental Forestry Studies and the Rutgers University Research Council.

Literature cited

- Agricultural Handbook No. 271. Silvics of Forest Trees of the United States. U.S. Dept. Agric. Forest Service, Washington, D.C.
- Ambler, M.A. 1965. Seven alien plant species. William L. Hutcheson Memorial Forest Bulletin 2(2):1-8.
- Baird, J. 1956. The ecology of the Watchung Reservation, Union Co., New Jersey. Publ. Dept. Botany Rutgers Univ., New Brunswick, N.J.
- Bard, G.E. 1952. Secondary succession on the Piedmont of New Jersey. Ecol. Monogr. 22:195-215.
- Biel, E.R. 1958. The climate of New Jersey. In *Economic Botany of New Jersey*. Rutgers Univ. Press, New Brunswick, N.J. Pp. 53-98.
- Braun, E. L. 1950. *Deciduous Forests of Eastern North America*. The Blakiston Co., Philadelphia, Pa.
- Bray, J.R., and J.T. Curtis. 1957. An ordination of the upland forest communities of southern Wisconsin. Ecol. Monogr. 27:325-349.
- Buell, M.F. 1957. The mature oak forest of Mettler Woods. William L. Hutcheson Memorial Forest Bulletin 1(1):16-19.
- _____, H.F. Buell and J.A. Small. 1954. Fire and the history of Mettler's Woods. Bull. Torrey Bot. Club 81:253-255.
- _____, A.N. Langford, D.W. Davidson and L.F. Ohmann. 1966. The upland forest continuum in northern New Jersey. Ecol. 47:417-432.
- _____, and W.A. Wistendahl. 1955. Flood plain forests of the Raritan River. Bull. Torrey Bot. Club 82:463-472.

- Cantlon, J.E. 1953. Vegetation and microclimates on north and south slopes of Cushetunk Mountain, New Jersey. *Ecol. Monogr.* 23:241-270.
- Fernald, M.L. 1950. *Gray's Manual of Botany*. Amer. Book Co. New York. 1632 p.
- Frei, K.R., and D.E. Fairbrothers. 1963. Floristic Study of the William L. Hutcheson Memorial Forest (New Jersey). *Bull. Torrey Bot. Club* 90:338-355.
- Galli, A.E. 1973. Bird species diversity and forest size on the New Jersey Piedmont. M.S. thesis. Rutgers University. New Brunswick, N.J.
- Gleason, H.A. 1962. *Plants of the Vicinity of New York*. Hafner Publ. Co. New York. 307 p.
- Harlow, W.M. 1959. *Trees of the Eastern and Central United States and Canada*. Dover Publ. Co. New York. 288 p.
- Horn, H.S. 1971. *The Adaptive Geometry of Trees*. Princeton Univ. Press. Princeton, N.J. 114 p.
- _____. 1975. Markovian properties of forest succession. In *Ecology and Evolution of Communities*. Eds: J.M. Diamond and M.J. Cody. Harvard Univ. Press. Cambridge, Mass. (in press).
- Kramer, R.J. 1971. Herrontown Woods. Stony Brook-Millstone Watershed Assn., Inc. Pennington, N.J. 44 p.
- Lewis, J.V., and H.B. Kummel. 1915. *The Geology of New Jersey*. Geol. Survey Bull. 15. Washington, D.C.
- McDonough, W.T., and M.F. Buell. 1956. The vegetation of Voorhees State Park, New Jersey. *Amer. Midl. Nat.* 56:473-490.
- Monk, C.D. 1957. Plant communities of Hutcheson Memorial Forest based on shrub distribution. *Bull. Torrey Bot. Club* 84:198-206.
- _____. 1961a. The vegetation of the William L. Hutcheson Memorial Forest, New Jersey. *Bull. Torrey Bot. Club* 88:156-166.
- _____. 1961b. Past and present influences on reproduction in the William L. Hutcheson Memorial Forest, New Jersey. *Bull. Torrey Bot. Club* 86:167-175.
- Reiners, N.M., and W.A. Reiners. 1965. Natural harvesting of trees. *William L. Hutcheson Memorial Forest Bulletin* 2(2):9-17.
- Robichaud, B., and M.F. Buell. 1973. *Vegetation of New Jersey. A Study of Landscape Diversity*. Rutgers Univ. Press, New Brunswick, N.J. 340 p.
- Small, J.A. 1973. Publications involving studies at the William L. Hutcheson Memorial Forest. *William L. Hutcheson Memorial Forest Bulletin* 3(1):27-29.
- Sulser, J.S. 1971. Twenty years of change in the Hutcheson Memorial Forest. *William L. Hutcheson Memorial Forest Bulletin* 2(4):15-25.
- Tedrow, J.C.F. 1963. *New Jersey Soils*. Rutgers College of Agriculture. Circular No. 601. Rutgers Univ. New Brunswick, N.J.
- Ugolini, F.C. 1964. Soil development on the red beds of New Jersey. *William L. Hutcheson Memorial Forest Bulletin* 2(1):1-34.
- U.S. Department of Commerce. 1959. *Climates of the States*. New Jersey. U.S. Weather Bureau. Gov. Printing Off. Washington, D.C.
- U.S. Soil Conservation Service. 1958-1967. *Cooperative Soil Surveys*. Somerset, Hunterdon and Mercer Counties. U.S. Dept. Agri. Soil Conservation Service and Rutgers Univ. Agric. Exp. Sta. Gov. Printing Off. Washington, D.C.
- Van Vechten, G.W., III, and M.F. Buell. 1959. The flood plain vegetation of the Millstone River, New Jersey. *Bull. Torrey Bot. Club* 86:219-227.
- Wales, B.A. 1972. Vegetation analysis of north and south edges in a mature oak-hickory forest. *Ecol. Monogr.* 42:451-471.
- Widmer, K. 1964. *The Geology and Geography of New Jersey*. Van Nostrand co. Princeton, N.J.
- Wistendahl, W.A. 1958. The flood plain of the Raritan River, New Jersey. *Ecol. Monogr.* 28:129-153.